



Entresto[®]
(sacubitril/valsartan) tablets
24/26 mg • 49/51 mg • 97/103 mg

for Heart Failure



Not actual patients.

WITH A HEALTHIER HEART, YOU CAN KEEP ON DOING WHAT YOU LOVE

ENTRESTO[®] is the #1 Heart Failure brand prescribed by cardiologists

What is ENTRESTO?

ENTRESTO is a prescription medicine used to treat adults with long-lasting (chronic) heart failure to help reduce the risk of death and hospitalization. ENTRESTO works better when the heart cannot pump a normal amount of blood to the body.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ENTRESTO?

ENTRESTO can harm or cause death to your unborn baby. Talk to your doctor about other ways to treat heart failure if you plan to become pregnant. If you get pregnant during treatment with ENTRESTO, tell your doctor right away.

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

TREATING YOUR HEART FAILURE MAY LEAD TO A WORLD OF POSSIBILITIES



Not actual patients.

IMPORTANT SAFETY INFORMATION

Do not take ENTRESTO if you:

- are allergic to any of the ingredients in ENTRESTO
- have had an allergic reaction including swelling of your face, lips, tongue, throat (angioedema) or trouble breathing while taking a type of medicine called an angiotensin-converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB)

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.



PROVEN SUPERIOR AT HELPING PEOPLE STAY ALIVE AND OUT OF THE HOSPITAL

In a study where the patients' hearts were stretched and weak, ENTRESTO[®] was proven superior at helping people stay alive and out of the hospital vs enalapril.

Heart Failure is not a one-time event

It's a long-lasting (chronic) condition that can get worse over time. ENTRESTO is the **#1 Heart Failure brand** prescribed by cardiologists and has helped over 1 million people with Heart Failure.

Try a month of ENTRESTO for free to see if it's right for you*

*Available to all patients when starting treatment, regardless of insurance coverage.

**Don't wait.
Talk to your doctor about ENTRESTO.**

Learn more at [ENTRESTO.com](https://www.entresto.com)

IMPORTANT SAFETY INFORMATION

Do not take ENTRESTO if you (cont):

- take an ACE inhibitor medicine. Do not take ENTRESTO for at least 36 hours before or after you take an ACE inhibitor medicine. Talk with your doctor or pharmacist before taking ENTRESTO if you are not sure if you take an ACE inhibitor medicine
- have diabetes and take a medicine that contains aliskiren

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

HOW ENTRESTO® WORKS



Entresto®
(sacubitril/valsartan) tablets
24/26 mg • 49/51 mg • 97/103 mg


ENTRESTO is a medicine specifically designed for Heart Failure. It contains 2 active ingredients that work in different ways.

Valsartan:

- Has been used for years to treat Heart Failure
- Reduces blood vessel tightening and the buildup of sodium and fluid

Sacubitril:

- Works unlike any other Heart Failure treatment and can only be found in ENTRESTO
- Helps relax blood vessels and decrease sodium and fluid in the body



Heart Failure can change the structure of your heart so it may not work as well.

ENTRESTO is the only Heart Failure medicine of its kind. It helps improve your heart's ability to pump blood to the body.

[Click here](#) to see how ENTRESTO works

IMPORTANT SAFETY INFORMATION

Before taking ENTRESTO tell your doctor about all of your medical conditions, including if you:

- have a history of hereditary angioedema
 - have kidney or liver problems
 - are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed.
- You should either take ENTRESTO or breastfeed. You should not do both

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

HOW TO TAKE ENTRESTO



Take your dose of ENTRESTO twice daily, exactly as prescribed. Your doctor will decide which dose is right for you.



24/26 mg



49/51 mg



97/103 mg

Pills shown not actual size.

Take ENTRESTO with or without food.

Take ENTRESTO at the same time each day to help you avoid missing a dose.

If you have been prescribed ENTRESTO and are currently taking an ACE inhibitor*:



Stop taking your ACE inhibitor*



Wait for 36 hours



Start taking ENTRESTO, as prescribed

*Such as enalapril or lisinopril.

IMPORTANT SAFETY INFORMATION

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you take potassium supplements or a salt substitute; nonsteroidal anti-inflammatory drugs (NSAIDs); lithium; or other medicines for high blood pressure or heart problems such as an ACE inhibitor, ARB, or aliskiren.

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

KNOW THE SYMPTOMS OF HEART FAILURE

Some symptoms of Heart Failure may go unnoticed or be mistaken for normal signs of aging. Knowing the difference can help you better manage your condition.

Symptoms of Heart Failure to watch out for:



Shortness of breath



Swelling



Feeling tired, fatigued



Coughing



Confusion



Rapid weight gain

Not actual patients.

Track your symptoms

Write your symptoms down each day and share them with your doctor. Call your doctor right away if you notice any new symptoms or if old symptoms have come back or gotten worse, as they could lead to Heart Failure hospitalization.

Start by using the ***Symptom Tracker*** on page 8.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of ENTRESTO?

ENTRESTO may cause serious side effects including:

- **Swelling of your face, lips, tongue and throat (angioedema) that may cause trouble breathing and death.** Get emergency medical help right away if you have symptoms of angioedema or trouble breathing. Do not take ENTRESTO again if you have had angioedema while taking ENTRESTO
- People who are Black or who have had angioedema and take ENTRESTO may have a higher risk of having angioedema

Please **[click here](#)** for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

TIPS TO HELP YOU MANAGE HEART FAILURE

In addition to your medication, keep in mind that a healthy diet and exercise are also important to manage your heart's health.



Eat a heart-healthy diet

Try to stay away from foods that are high in salt, as too much can make swelling worse.



Get moving

Even a little bit of activity can help. Just be sure to talk to your doctor first about which activities are good for you.



Weigh yourself regularly

Check your weight every morning and write it down in your Symptom Tracker. If you notice you've gained 3 or more pounds in 1 day, call your doctor right away.



Be open with your doctor

Tell your doctor what's important to you. It will help them come up with the best management plan for you.

Start by using the ***Medicine Tracker*** on page 9.

IMPORTANT SAFETY INFORMATION

ENTRESTO may cause serious side effects including (cont):

- **low blood pressure (hypotension)**, which may be more common if you take water pills. Call your doctor if you become dizzy or lightheaded, or you develop extreme fatigue
- **kidney problems**
- **increased amount of potassium in your blood (hyperkalemia)**

The most common side effects were low blood pressure, high potassium, cough, dizziness, and kidney problems.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

SYMPTOM TRACKER

Record your symptoms every day.

Print more pages or download a copy to take to your next doctor's appointment [here](#).



Entresto®
(sacubitril/valsartan) tablets
24/26mg • 49/51mg • 97/103mg

DATE: _____

 WEIGHT	<p>Weigh yourself each morning at the same time. If you notice you've gained 3 or more pounds in 1 day, call your doctor right away.</p> <p style="text-align: right;">_____ LBS.</p>				
 FEELING TIRED	NONE	BETTER	SAME	WORSE	NEW
 SWELLING	NONE	BETTER	SAME	WORSE	NEW
 COUGHING	NONE	BETTER	SAME	WORSE	NEW
 SHORTNESS OF BREATH	NONE	BETTER	SAME	WORSE	NEW

DATE: _____

 WEIGHT	<p>Weigh yourself each morning at the same time. If you notice you've gained 3 or more pounds in 1 day, call your doctor right away.</p> <p style="text-align: right;">_____ LBS.</p>				
 FEELING TIRED	NONE	BETTER	SAME	WORSE	NEW
 SWELLING	NONE	BETTER	SAME	WORSE	NEW
 COUGHING	NONE	BETTER	SAME	WORSE	NEW
 SHORTNESS OF BREATH	NONE	BETTER	SAME	WORSE	NEW

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

TRACK YOUR TREATMENTS

It's important to know why each medicine has been prescribed, what it does, and when to take it. This includes over-the-counter products, vitamins, and dietary supplements. Print out as many pages as you need to take to your next doctor's appointment.

MEDICINE _____ CONDITION _____
DOSE/HOW OFTEN _____
PRESCRIBING DOCTOR _____ DATE _____

MEDICINE _____ CONDITION _____
DOSE/HOW OFTEN _____
PRESCRIBING DOCTOR _____ DATE _____

MEDICINE _____ CONDITION _____
DOSE/HOW OFTEN _____
PRESCRIBING DOCTOR _____ DATE _____

MEDICINE _____ CONDITION _____
DOSE/HOW OFTEN _____
PRESCRIBING DOCTOR _____ DATE _____

MEDICINE _____ CONDITION _____
DOSE/HOW OFTEN _____
PRESCRIBING DOCTOR _____ DATE _____

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

IMPORTANT FACTS ABOUT ENTRESTO®

What is the most important information I should know about ENTRESTO?

ENTRESTO can harm or cause death to your unborn baby. Talk to your doctor about other ways to treat heart failure if you plan to become pregnant. If you get pregnant during treatment with ENTRESTO, tell your doctor right away.

WHAT IS ENTRESTO?

ENTRESTO is a prescription medicine used to treat adults with long-lasting (chronic) heart failure to help reduce the risk of death and hospitalization. ENTRESTO works better when the heart cannot pump a normal amount of blood to the body. It is also used to treat certain children 1 year of age and older who have symptomatic heart failure.

It is not known if ENTRESTO is safe and effective in children under 1 year of age.

Do not take ENTRESTO if you:

- are allergic to any of the ingredients in ENTRESTO.
- have had an allergic reaction including swelling of your face, lips, tongue, throat, or trouble breathing while taking a type of medicine called an angiotensin-converting enzyme (ACE) inhibitor or angiotensin II receptor blocker (ARB).
- take an ACE inhibitor medicine. **Do not take ENTRESTO for at least 36 hours before or after you take an ACE inhibitor medicine.** Talk with your doctor or pharmacist before taking ENTRESTO if you are not sure if you take an ACE inhibitor medicine.
- have diabetes and take a medicine that contains aliskiren.

Before you take ENTRESTO, tell your doctor about all of your medical conditions, including if you:

- have a history of hereditary angioedema
- have kidney or liver problems
- are pregnant or plan to become pregnant. See **“What is the most important information I should know about ENTRESTO?”**
- are breastfeeding or plan to breastfeed. It is not known if ENTRESTO passes into your breast milk. You and your doctor should decide if you will take ENTRESTO or breastfeed. You should not do both.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using ENTRESTO with certain other medicines may affect each other. Using ENTRESTO with other medicines can cause serious side effects. Especially tell your doctor if you take: potassium supplements or a salt substitute; nonsteroidal anti-inflammatory drugs (NSAIDs); lithium; other medicines for high blood pressure or heart problems such as an ACE inhibitor, ARB, or aliskiren. Keep a list of your medicines to show your doctor and pharmacist when you get a new medicine.

Please [click here](#) for full Prescribing Information, including Boxed WARNING.



Entresto[®]
(sacubitril/valsartan) tablets
24/26mg · 49/51mg · 97/103mg

What are the possible side effects of ENTRESTO?

ENTRESTO may cause serious side effects including:

- See “**What is the most important information I should know about ENTRESTO?**”
- **Serious allergic reactions causing swelling of your face, lips, tongue, and throat (angioedema) that may cause trouble breathing and death.** Get emergency medical help right away if you have symptoms of angioedema or trouble breathing. Do not take ENTRESTO again if you have had angioedema during treatment with ENTRESTO.
- People who are Black and take ENTRESTO may have a higher risk of having angioedema than people who are not Black and take ENTRESTO.
- People who have had angioedema before taking ENTRESTO may have a higher risk of having angioedema than people who have not had angioedema before taking ENTRESTO. See “**Who should not take ENTRESTO?**”
- **Low blood pressure (hypotension).** Low blood pressure may be more common if you also take water pills. Call your doctor if you become dizzy or lightheaded, or you develop extreme fatigue.
- **Kidney problems.** Your doctor will check your kidney function during your treatment with ENTRESTO. If you have changes in your kidney function tests, you may need a lower dose of ENTRESTO or may need to stop taking ENTRESTO for a period of time.
- **Increased amount of potassium in your blood (hyperkalemia).** Your doctor will check your potassium blood level during your treatment with ENTRESTO.

These are not all the possible side effects of ENTRESTO. Call your doctor for medical advice about side effects.

The most common side effects were low blood pressure, high potassium, cough, dizziness, and kidney problems.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

This information is not comprehensive. To learn more, talk to your health care provider or pharmacist, visit www.entresto.com to obtain the FDA-approved product labeling, or call 1-888-ENTRESTO.

Please [click here](#) for full Prescribing Information, including Boxed WARNING.

ENTRESTO® OFFERS SAVINGS AND SUPPORT FROM THE START

Enspire
PROGRAM
from Entresto®
SUPPORT DESIGNED WITH YOU IN MIND

We're committed to helping make ENTRESTO more affordable for everyone—it's covered by most insurance plans and has the lowest branded co-pay for over 99% of people with Medicare.^{1*}

Try a month of
ENTRESTO for free to
see if it's right for you^{†‡}

Fill your prescription
for as little as \$10[‡]
For eligible commercially insured patients.[§]
Pay as little as \$10 for a 30-, 60-, or 90-day supply.

The **ENSPIRE Program from ENTRESTO®** offers support designed with you in mind. It provides tips and tools that can help you during treatment. **You can:**

- Improve your diet with healthy recipes, meal plans, and how-to guides, and get activity tips and resources
- Learn how to set up a medicine schedule, get tips for staying on track, and find information on how others can help

Visit [ENTRESTO.com/cost](https://www.entresto.com/cost) or call **1-888-ENTRESTO** to learn more about lifestyle, treatment, and financial support.

 facebook.com/entresto

 [entresto_sacubitril_valsartan](https://www.instagram.com/entresto_sacubitril_valsartan)

 [YouTube.com/LivingWithHF](https://www.youtube.com/LivingWithHF)

[Click here to learn more](#)


Entresto®
(sacubitril/valsartan) tablets
24/26mg • 49/51mg • 97/103mg

*Out-of-pocket costs will vary from plan to plan. See your plan's coverage information for more details.

†Available to all patients when starting treatment, regardless of insurance coverage. One per patient.

‡Limitations apply. See Terms and Conditions at [ENTRESTO.com/cost](https://www.entresto.com/cost).

§Not valid under Medicare, Medicaid, or any other federal or state program.

Reference:

1. Data on file. Novartis Pharmaceuticals Corp; 2021.

Please [click here](#) for full Prescribing Information, including Boxed WARNING, and see Important Facts About ENTRESTO on pages 10 and 11.

ENTRESTO and the ENTRESTO logo are registered trademarks of Novartis AG.

 **NOVARTIS**

Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936-1080

© 2023 Novartis

Printed in USA

5/23

280927